

Centers of symmetric cellular algebras *

Yanbo Li

Department of Information and Computing Sciences,
Northeastern University at Qinhuangdao;
Qinhuangdao, 066004, P.R. China

School of Mathematics Sciences, Beijing Normal University;
Beijing, 100875, P.R. China
E-mail: liyanbo707@163.com

November 18, 2009

Abstract

Let R be an integral domain and A a symmetric cellular algebra over R with a cellular basis $\{C_{S,T}^\lambda \mid \lambda \in \Lambda, S, T \in M(\lambda)\}$. We will construct an ideal $L(A)$ of the center of A and prove that $L(A)$ contains the so-called Higman ideal. When R is a field, we prove that the dimension of $L(A)$ is not less than the number of non-isomorphic simple A -modules.

1 Introduction

In 1996, Graham and Lehrer [8] introduced cellular algebras in order to provide a systematic framework for studying the representation theory of a class of algebras. By the theory of cellular algebras, one can parameterize simple modules for a finite dimensional cellular algebra by methods in linear algebra. Many classes of algebras from mathematics and physics are found to be cellular, including Hecke algebras of finite type, Ariki-Koike algebras, q -Schur algebras, Brauer algebras, Temperley-Lieb algebras, cyclotomic Temperley-Lieb algebras, partition algebras, Birman-Wenzl algebras and so on, see [7], [8], [14], [15], [16] for details.

*keywords: symmetric cellular algebras, center.

There are many papers on centers of Hecke algebras of finite type, which are all cellular algebras [7]. In [11], Jones found bases for centers of Hecke algebras of type A over $\mathbb{Q}[q, q^{-1}]$, where q is an indeterminant. This basis is an analog of conjugacy class sum in a group algebra. In [10], Geck and Rouquier found bases for the centers of generic Hecke algebras over $\mathbb{Z}[q, q^{-1}]$ with q an indeterminant. However, it is not easy to write the basis explicitly. Then one should ask, is there any basis which can be written explicitly? In [3], Francis gave an integral minimal basis for the center of a Hecke algebra. Then in [4], he used the minimal basis approach to provide a way of describing and calculating elements of the minimal basis for the center of an Iwahori-Hecke algebra which is entirely combinatorial. In [6], Francis and Jones found an explicit non-recursive expression for the coefficients appearing in these linear combinations for the Hecke algebras of type A. The relations between the so-called Jucys-Murphy elements and centers of Hecke algebras are also investigated widely. In [2], Dipper and James conjectured that the center of a Hecke algebra of type A consists of symmetric polynomials in the Jucys-Murphy elements. This conjecture was proved by Francis and Graham [5] in 2006. An analogous conjecture for Ariki-Koike algebras is still open.

The fact that Hecke algebras of finite type are all cellular leads us to considering how to describe the centers of Hecke algebras by cellular bases. Furthermore, how to describe the center of a cellular algebra in general? Clearly, most of the approaches for studying Hecke algebras can not be used directly for cellular algebras, since we have no Weyl group structure to use. Then we must look for some new method. In fact, the symmetry of Hecke algebras provides us a way. We will do some work on the centers of symmetric cellular algebras in this paper.

In order to describe our result exactly, we fix some notations first. Let A be a symmetric cellular R -algebra with a non-degenerate symmetric bilinear form $f : A \times A \rightarrow R$. Then f determines a map $\tau : A \rightarrow R$ which is defined by $\tau(a) = f(a, 1)$ for every $a \in A$. We call τ a symmetrizing trace. Denote by $\{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ the dual basis determined by τ . Let $H(A) = \{ \sum_{\lambda \in \Lambda, S, T \in M(\lambda)} C_{S,T}^\lambda a D_{S,T}^\lambda \mid a \in A \}$. It is the Higman ideal of $Z(A)$. For any $\lambda \in \Lambda$ and some $T \in M(\lambda)$, set $x_\lambda = \sum_{S \in M(\lambda)} C_{S,T}^\lambda D_{S,T}^\lambda$, where x_λ is independent of T , and $L(A) = \{ \sum_{\lambda \in \Lambda} r_\lambda x_\lambda \mid r_\lambda \in R \}$. Then the main result of this paper is as follows.

Theorem. *Let A be a symmetric cellular algebra with a cellular basis $\{C_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ and the dual basis $\{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ determined by a symmetrizing trace τ . Then*

- (1) $L(A)$ is an ideal of $Z(A)$ and contains the Higman ideal.
- (2) $L(A)$ is independent of the choice of τ .
- (3) If R is a field, then the dimension of $L(A)$ is not less than the number of non-

isomorphic simple A -modules.

This theorem enlarges the well known Higman ideal to a new one for the center of a symmetric cellular algebra.

2 Preliminaries

In this section, we first recall some basic results on symmetric algebras and cellular algebras, which is needed in the following sections. The so-called Higman ideal is also described. References for this section are the books [1] and [9].

Let R be a commutative ring with identity and A an associative R -algebra. As an R -module, A is finitely generated and free. Suppose that there exists an R -bilinear map $f : A \times A \rightarrow R$. We say that f is non-degenerate if the determinant of the matrix $(f(a_i, a_j))_{a_i, a_j \in B}$ is a unit in R for some R -basis B of A . We say f is associative if $f(ab, c) = f(a, bc)$ for all $a, b, c \in A$, and symmetric if $f(a, b) = f(b, a)$ for all $a, b \in A$.

Definition 2.1. *An R -algebra A is called symmetric if there is a non-degenerate associative symmetric bilinear form f on A . Define an R -linear map $\tau : A \rightarrow R$ by $\tau(a) = f(a, 1)$. We call τ a symmetrizing trace.*

Let A be a symmetric algebra with a basis $B = \{a_i \mid i = 1, \dots, n\}$ and τ a symmetrizing trace. Denote by $D = \{D_i \mid i = 1, \dots, n\}$ the basis determined by the requirement that $\tau(D_j a_i) = \delta_{ij}$ for all $i, j = 1, \dots, n$. We will call D the dual basis of B . For arbitrary $1 \leq i, j \leq n$, we write $a_i a_j = \sum_k r_{ijk} a_k$, where $r_{ijk} \in R$. Fix a τ for A . Then in [13], we proved the following lemma.

Lemma 2.2. *Let A be a symmetric algebra with a basis B and the dual basis D . Then the following hold: $a_i D_j = \sum_k r_{kij} D_k$; $D_i a_j = \sum_k r_{jki} D_k$. $\square$*

We now consider the set $\{\sum_i D_i a a_i \mid a \in A\}$. It is well known that $\{\sum_i D_i a a_i \mid a \in A\}$ is an ideal of the center of A , see [1]. We here give a direct proof by the lemma above.

Proposition 2.3. *Let A be a symmetric algebra with a basis B and the dual basis D . Then $\{\sum_i D_i a a_i \mid a \in A\}$ is an ideal of the center of A .*

Proof: For arbitrary $a_j \in B$ and $a \in A$, we have

$$\sum_i D_i a a_i a_j = \sum_{i,k} r_{ijk} D_i a a_k,$$

and

$$\sum_i a_j D_i a a_i = \sum_{i,k} r_{kji} D_k a a_i.$$

Obviously, the right sides of the above two equations are equal. Then $\{\sum_i D_i a a_i \mid a \in A\} \subseteq Z(A)$. It is clear that the set is an ideal of the center of A . $\square$

The ideal $H(A) = \{\sum_i D_i a a_i \mid a \in A\}$ is called Higman ideal of the center of the algebra A . It is independent the choice of the dual bases.

The following proposition is also proved in [13].

Proposition 2.4. *Suppose that A is a symmetric R -algebra with a basis $\{a_i \mid i = 1, \dots, n\}$. Let τ, τ' be two symmetrizing traces. Denote by $\{D_i \mid i = 1, \dots, n\}$ and $\{D'_i \mid i = 1, \dots, n\}$ the dual bases determined by τ and τ' respectively. Then for any $1 \leq i \leq n$, we have $D'_i = \sum_{j=1}^n \tau(a_j D'_i) D_j$.* $\square$

We now recall the definition of cellular algebras introduced by Graham and Lehrer [8] and some well known results.

Definition 2.5. ([8] 1.1) *Let R be a commutative ring with identity. An associative unital R -algebra is called a cellular algebra with cell datum (Λ, M, C, i) if the following conditions are satisfied:*

(C1) *The finite set Λ is a poset. Associated with each $\lambda \in \Lambda$, there is a finite set $M(\lambda)$. The algebra A has an R -basis $\{C_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$.*

(C2) *The map i is an R -linear anti-automorphism of A with $i^2 = \text{id}$ which sends $C_{S,T}^\lambda$ to $C_{T,S}^\lambda$.*

(C3) *If $\lambda \in \Lambda$ and $S, T \in M(\lambda)$, then for any element $a \in A$, we have*

$$a C_{S,T}^\lambda \equiv \sum_{S' \in M(\lambda)} r_a(S', S) C_{S',T}^\lambda \pmod{A(< \lambda)},$$

where $r_a(S', S) \in R$ is independent of T and where $A(< \lambda)$ is the R -submodule of A generated by $\{C_{S'',T''}^\mu \mid S'', T'' \in M(\mu), \mu < \lambda\}$.

Apply i to the equation in (C3), we obtain

$$(C3') \quad C_{T,S}^\lambda i(a) \equiv \sum_{S' \in M(\lambda)} r_a(S', S) C_{T,S'}^\lambda \pmod{A(< \lambda)}.$$

By Definition 2.5, it is easy to check that

$$C_{S,S}^\lambda C_{T,T}^\lambda \equiv \Phi(S, T) C_{S,T}^\lambda \pmod{A(< \lambda)},$$

where $\Phi(S, T) \in R$ depends only on S and T .

Let A be a cellular algebra with cell datum (Λ, M, C, i) . We recall the definition of cell modules.

Definition 2.6. ([8] 2.1) For each $\lambda \in \Lambda$, define the left A -module $W(\lambda)$ as follows: $W(\lambda)$ is a free R -module with basis $\{C_S \mid S \in M(\lambda)\}$ and A -action defined by

$$aC_S = \sum_{S' \in M(\lambda)} r_a(S', S)C_{S'} \quad (a \in A, S \in M(\lambda)),$$

where $r_a(S', S)$ is the element of R defined in (C3).

For a cell module $W(\lambda)$, define a bilinear form $\Phi_\lambda : W(\lambda) \times W(\lambda) \longrightarrow R$ by $\Phi_\lambda(C_S, C_T) = \Phi(S, T)$, extended bilinearly and define

$$\text{rad}(\lambda) := \{x \in W(\lambda) \mid \Phi_\lambda(x, y) = 0 \text{ for all } y \in W(\lambda)\}.$$

Then Graham and Lehrer proved the following results in [8].

Theorem 2.7. [8] Let K be a field and A a finite dimensional cellular algebra. Denote the A -module $W(\lambda)/\text{rad} \lambda$ by L_λ , where $\lambda \in \Lambda$ with $\Phi_\lambda \neq 0$. Let $\Lambda_0 = \{\lambda \in \Lambda \mid \Phi_\lambda \neq 0\}$. Then the set $\{L_\lambda \mid \lambda \in \Lambda_0\}$ is a complete set of (representative of equivalence classes of) absolutely simple A -modules. $\square$

3 Centers of symmetric cellular algebras

Let A be a symmetric cellular algebra with a cell datum (Λ, M, C, i) . Denote the dual basis by $D = \{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$, which satisfies

$$\tau(C_{S,T}^\lambda D_{U,V}^\mu) = \begin{cases} 1, & \lambda = \mu, S = U, T = V; \\ 0, & \text{otherwise.} \end{cases}$$

For any $\lambda, \mu \in \Lambda, S, T \in M(\lambda), U, V \in M(\mu)$, write

$$C_{S,T}^\lambda C_{U,V}^\mu = \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} r_{(S,T,\lambda),(U,V,\mu),(X,Y,\varepsilon)} C_{X,Y}^\varepsilon.$$

Then in [13], we proved the following lemma.

Lemma 3.1. Let A be a symmetric cellular algebra with a basis B . Let D be the dual basis determined by a given τ . For arbitrary $\lambda, \mu \in \Lambda$ and $S, T, P, Q \in M(\lambda), U, V \in M(\mu)$, the following hold:

- (1) $D_{U,V}^\mu C_{S,T}^\lambda = \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} r_{(S,T,\lambda),(X,Y,\varepsilon),(U,V,\mu)} D_{X,Y}^\varepsilon.$
- (2) $C_{S,T}^\lambda D_{U,V}^\mu = \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} r_{(X,Y,\varepsilon),(S,T,\lambda),(U,V,\mu)} D_{X,Y}^\varepsilon.$

- (3) $C_{S,T}^\lambda D_{S,T}^\lambda = C_{S,P}^\lambda D_{S,P}^\lambda$. (4) $D_{S,T}^\lambda C_{S,T}^\lambda = D_{P,T}^\lambda C_{P,T}^\lambda$.
 (5) $C_{S,T}^\lambda D_{P,Q}^\lambda = 0$ if $T \neq Q$. (6) $D_{P,Q}^\lambda C_{S,T}^\lambda = 0$ if $P \neq S$.
 (7) $C_{S,T}^\lambda D_{U,V}^\mu = 0$ if $\mu \not\leq \lambda$. (8) $D_{U,V}^\mu C_{S,T}^\lambda = 0$ if $\mu \not\leq \lambda$. $\square$

Let A be a symmetric cellular R -algebra with a symmetrizing trace τ . The dual basis $\{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ is determined by τ . Then the Higman ideal is $H(A) = \left\{ \sum_{\lambda \in \Lambda, S, T \in M(\lambda)} C_{S,T}^\lambda a D_{S,T}^\lambda \mid a \in A \right\}$. For any $\lambda \in \Lambda$ and $T \in M(\lambda)$, set $x_\lambda = \sum_{S \in M(\lambda)} C_{S,T}^\lambda D_{S,T}^\lambda$ and $L(A) = \left\{ \sum_{\lambda \in \Lambda} r_\lambda x_\lambda \mid r_\lambda \in R \right\}$. Now we are in a position to give the main result of this paper.

Theorem 3.2. *Let A be a symmetric cellular algebra with a cellular basis $\{C_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ and the dual basis $\{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ determined by a symmetrizing trace τ . Then*

- (1) $L(A)$ is an ideal of $Z(A)$ and contains the Higman ideal $H(A)$.
 (2) $L(A)$ is independent of the choice of τ .
 (3) If R is a field, then the dimension of $L(A)$ is not less than the number of non-isomorphic simple A -modules.

Proof: (1) Firstly, we show that $H(A) \subseteq L(A)$.

Clearly, we only need to show that $l := \sum_{S, T \in M(\lambda), \lambda \in \Lambda} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda \in L(A)$ for any $C_{U,V}^\mu \in B$, where $\mu \in \Lambda, U, V \in M(\mu)$. We divide l into three parts: $l = l_{\lambda=\mu} + l_{\lambda < \mu} + l_{\lambda \not\leq \mu}$, where

$$l_{\lambda=\mu} := \sum_{S, T \in M(\lambda), \lambda=\mu} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda$$

and the other two parts are defined similarly.

By Lemma 3.1 (7), $l_{\lambda \not\leq \mu} = 0$.

We now show that

$$l_{\lambda=\mu} = \Phi_\mu(C_U, C_V) x_\mu.$$

By Lemma 3.1 (5), $C_{U,V}^\mu D_{X,Y}^\mu = 0$ if $V \neq Y$. Then $l_{\lambda=\mu} = \sum_{X \in M(\mu)} C_{X,V}^\mu C_{U,V}^\mu D_{X,V}^\mu$.

By Definition 2.5, we have

$$l_{\lambda=\mu} = \sum_{X \in M(\mu)} \Phi_\mu(C_U, C_V) C_{X,V}^\mu D_{X,V}^\mu + \sum_{\eta < \mu, P, Q \in M(\eta)} r_{P,Q,\eta} C_{P,Q}^\eta D_{X,V}^\mu,$$

where $r_{P,Q,\eta} \in R$. Note that by Lemma 3.1 (7), $\sum_{\eta < \mu, P, Q \in M(\mu)} r_{P,Q,\eta} C_{P,Q}^\eta D_{X,V}^\mu = 0$, then $l_{\lambda=\mu} = \Phi_\mu(C_U, C_V) x_\mu$. This implies that $l_{\lambda=\mu} \in L(A)$.

Now let us consider $l_{\lambda < \mu}$. For arbitrary $\lambda < \mu$, we show that

$$\sum_{S,T \in M(\lambda)} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda = \sum_{T \in M(\lambda)} r_{(S,T,\lambda),(U,V,\mu),(S,T,\lambda)} x_\lambda.$$

Note that

$$\sum_{S,T \in M(\lambda)} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda = \sum_{S,T \in M(\lambda)} \left(\sum_{\varepsilon \in \Lambda, X,Y \in M(\varepsilon)} r_{(S,T,\lambda),(U,V,\mu),(X,Y,\varepsilon)} C_{X,Y}^\varepsilon \right) D_{S,T}^\lambda.$$

By (C3)' of Definition 2.5, if $\varepsilon \not\leq \lambda$, then $r_{(S,T,\lambda),(U,V,\mu),(X,Y,\varepsilon)} = 0$. By Lemma 3.1 (7), if $\varepsilon < \lambda$, then $C_{X,Y}^\varepsilon D_{S,T}^\lambda = 0$. Thus

$$\sum_{S,T \in M(\lambda)} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda = \sum_{S,T \in M(\lambda)} \sum_{X,Y \in M(\lambda)} r_{(S,T,\lambda),(U,V,\mu),(X,Y,\lambda)} C_{X,Y}^\lambda D_{S,T}^\lambda.$$

By (C3)' of Definition 2.5, if $X \neq S$, then $r_{(S,T,\lambda),(U,V,\mu),(X,Y,\lambda)} = 0$. By Lemma 3.1 (5), if $Y \neq T$, then $C_{X,Y}^\lambda D_{S,T}^\lambda = 0$. Hence,

$$\sum_{S,T \in M(\lambda)} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda = \sum_{S,T \in M(\lambda)} r_{(S,T,\lambda),(U,V,\mu),(S,T,\lambda)} C_{S,T}^\lambda D_{S,T}^\lambda.$$

Note that for arbitrary $S, S' \in M(\lambda)$, by (C3)' of Definition 2.5.

$$r_{(S,T,\lambda),(U,V,\mu),(S,T,\lambda)} = r_{(S',T,\lambda),(U,V,\mu),(S',T,\lambda)}.$$

We get

$$\sum_{S,T \in M(\lambda)} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda = \sum_{T \in M(\lambda)} r_{(S,T,\lambda),(U,V,\mu),(S,T,\lambda)} x_\lambda.$$

This implies $l_{\lambda < \mu} \in L(A)$. Then we obtain $l \in L(A)$.

Secondly, we show that $L(A) \subseteq Z(A)$.

We only need to show that $x_\lambda C_{U,V}^\mu = C_{U,V}^\mu x_\lambda$ for arbitrary $\lambda \in \Lambda$ and $\mu \in \Lambda, U, V \in M(\mu)$.

On one hand, by Lemma 3.1 (1),

$$\begin{aligned} x_\lambda C_{U,V}^\mu &= \sum_{S \in M(\lambda)} C_{S,T}^\lambda D_{S,T}^\lambda C_{U,V}^\mu \\ &= \sum_{S \in M(\lambda)} \sum_{\varepsilon \in \Lambda, X,Y \in M(\varepsilon)} r_{(U,V,\mu),(X,Y,\varepsilon),(S,T,\lambda)} C_{S,T}^\lambda D_{X,Y}^\varepsilon. \end{aligned}$$

By a similar method as in the first part, we get

$$x_\lambda C_{U,V}^\mu = \sum_{S,X \in M(\lambda)} r_{(U,V,\mu),(X,T,\lambda),(S,T,\lambda)} C_{S,T}^\lambda D_{X,T}^\lambda.$$

On the other hand,

$$\begin{aligned} C_{U,V}^\mu x_\lambda &= \sum_{S \in M(\lambda)} \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} r_{(U,V,\mu),(S,T,\lambda),(X,Y,\varepsilon)} C_{X,Y}^\varepsilon D_{S,T}^\lambda \\ &= \sum_{S, X \in M(\lambda)} r_{(U,V,\mu),(S,T,\lambda),(X,T,\lambda)} C_{X,T}^\lambda D_{S,T}^\lambda. \end{aligned}$$

So $x_\lambda C_{U,V}^\mu = C_{U,V}^\mu x_\lambda$ for arbitrary $\lambda, \mu \in \Lambda$, $U, V \in M(\mu)$, that is, $L(A) \subseteq Z(A)$.

Finally, we show that $L(A)$ is an ideal of $Z(A)$.

It suffices to show that for arbitrary $c \in Z(A)$ and $\lambda \in \Lambda$, the element $cx_\lambda \in L(A)$, that is, $\sum_{S \in M(\lambda)} C_{S,T}^\lambda c D_{S,T}^\lambda \in L(A)$.

Since c is R -linear combination of elements of B , then we only need to prove that for arbitrary $C_{U,V}^\mu \in B$, the element $\sum_{S \in M(\lambda)} C_{S,T}^\lambda C_{U,V}^\mu D_{S,T}^\lambda \in L(A)$. Clearly, this element is equal to

$$\sum_{S \in M(\lambda)} \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} r_{(S,T,\lambda),(U,V,\mu),(X,Y,\varepsilon)} C_{X,Y}^\varepsilon D_{S,T}^\lambda.$$

We know that it is equal to $r_{(S,T,\lambda),(U,V,\mu),(X,Y,\varepsilon)} x_\lambda$ by a similar way as in the first part. This implies that $\sum_{S \in M(\lambda)} C_{S,T}^\lambda c D_{S,T}^\lambda \in L(A)$.

(2) $L(A)$ is independent of the choice of τ .

Let τ, τ' be two non-equal symmetrizing traces and D, d the dual bases determined by τ and τ' respectively. For arbitrary $d_{S,T}^\lambda \in d$, by Proposition 2.4, we have

$$d_{S,T}^\lambda = \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} \tau(C_{X,Y}^\varepsilon d_{S,T}^\lambda) D_{X,Y}^\varepsilon.$$

Then by Lemma 3.1,

$$\begin{aligned} \sum_{S \in M(\lambda)} C_{S,T}^\lambda d_{S,T}^\lambda &= \sum_{S \in M(\lambda)} \sum_{\varepsilon \in \Lambda, X, Y \in M(\varepsilon)} \tau(C_{X,Y}^\varepsilon d_{S,T}^\lambda) C_{S,T}^\lambda D_{X,Y}^\varepsilon \\ &= \sum_{S \in M(\lambda)} \sum_{X \in M(\lambda)} \tau(C_{X,T}^\lambda d_{S,T}^\lambda) C_{S,T}^\lambda D_{X,T}^\lambda. \end{aligned}$$

By the definition of τ , we have $\tau(C_{X,T}^\lambda d_{S,T}^\lambda) = \tau(d_{S,T}^\lambda C_{X,T}^\lambda)$. Then by Lemma 3.1, $\tau(d_{S,T}^\lambda C_{X,T}^\lambda) = 0$ if $X \neq S$, that is,

$$\sum_{S \in M(\lambda)} C_{S,T}^\lambda d_{S,T}^\lambda = \sum_{S \in M(\lambda)} \tau(C_{S,T}^\lambda d_{S,T}^\lambda) C_{S,T}^\lambda D_{S,T}^\lambda.$$

We now need to show $\tau(C_{S,T}^\lambda d_{S,T}^\lambda)$ is independent of S . It is clear by the equations $d_{S,T}^\lambda C_{S,T}^\lambda = d_{S',T}^\lambda C_{S',T}^\lambda$ for arbitrary $S' \in M(\lambda)$.

(3) We only need to find $|\Lambda_0|$ R -linear independent elements in $L(A)$, where $|\Lambda_0|$ is the number of the elements in Λ_0 . By the definition of Λ_0 , for each $\lambda \in \Lambda_0$, there exist $S, T \in M(\lambda)$, such that $\Phi_\lambda(C_S, C_T) \neq 0$. Write $x_\lambda = \sum_{U \in M(\lambda)} C_{U,T}^\lambda D_{U,T}^\lambda$.

By Lemma 3.1, we know that the coefficient of $D_{S,T}^\lambda$ in the expansion of $C_{S,T}^\lambda D_{S,T}^\lambda$ is $r_{(S,T,\lambda),(S,T,\lambda),(S,T,\lambda)} = \Phi_\lambda(C_S, C_T) \neq 0$ and is 0 in the expansion of $C_{U,T}^\lambda D_{U,T}^\lambda$ for any $U \neq S$. That is, the coefficient of $D_{S,T}^\lambda$ in the expansion of x_λ is not zero. We also know that the coefficient of $D_{S,T}^\lambda$ in the expansion of x_μ is zero for any $\mu \not\leq \lambda$. Now let $\sum_{\lambda \in \Lambda_0} r_\lambda x_\lambda = 0$ and μ a minimal element in Λ_0 . Then r_μ must be zero. By induction, we know that $r_\lambda = 0$ for each $\lambda \in \Lambda_0$. This implies that $\{x_\lambda \mid \lambda \in \Lambda_0\}$ is R -linear independent. That is, $\dim_R L(A)$ is not less than the number of (representatives of equivalence classes of) simple A -modules. $\square$

By a similar way, we obtain the following result.

Theorem 3.3. Suppose that R is a commutative ring with identity and A a symmetric cellular algebra over R with a cellular basis $B = \{C_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$ and the dual basis $D = \{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$. Denote the set of the R -linear combination of the elements of the set $\{x'_\lambda = \sum_{T \in M(\lambda)} D_{S,T}^\lambda C_{S,T}^\lambda \mid \lambda \in \Lambda\}$ by $L(A)'$.

Then the following hold:

- (1) $L(A)'$ is an ideal of $Z(A)$ and contains the Higman ideal.
- (2) $L(A)'$ is independent of the choice of τ .
- (3) If R is a field, then the dimension of $L(A)'$ is not less than the number of (representatives of equivalence classes of) simple A -modules.

We now give some examples of $L(A)$.

Example Let K be a field and Q the following quiver

$$\begin{array}{ccccccc} \bullet & \xrightarrow{\alpha_1} & \bullet & \xrightarrow{\alpha_2} & \cdots & \bullet & \xrightarrow{\alpha_{n-1}} & \bullet \\ 1 & \xleftarrow{\alpha'_1} & 2 & \xleftarrow{\alpha'_2} & 3 & \cdots & n-1 & \xleftarrow{\alpha'_{n-1}} & n \end{array}$$

with relation ρ given as follows.

- (1) all paths of length ≥ 3 ;
 - (2) $\alpha'_i \alpha_i - \alpha_{i+1} \alpha'_{i+1}$, $i = 1, \dots, n-2$;
 - (3) $\alpha_i \alpha_{i+1}$, $\alpha'_{i+1} \alpha'_i$, $i = 1, \dots, n-2$.
- Let $A = K(Q, \rho)$. Define τ by
- (1) $\tau(e_1) = \dots = \tau(e_n) = 1$;

$$(2) \tau(\alpha_i \alpha'_i) = \tau(\alpha'_i \alpha_i) = 1, i = 1, \dots, n-1;$$

$$(3) \tau(\alpha_i) = \tau(\alpha'_i) = 0.$$

Then A is a symmetric cellular algebra with a cellular basis

$$e_1; \begin{matrix} \alpha_1 \alpha'_1 & \alpha_1 \\ \alpha'_1 & e_2 \end{matrix}; \begin{matrix} \alpha_2 \alpha'_2 & \alpha_2 \\ \alpha'_2 & e_3 \end{matrix}; \dots; \begin{matrix} \alpha_{n-1} \alpha'_{n-1} & \alpha_{n-1} \\ \alpha'_{n-1} & e_n \end{matrix}; \alpha'_{n-1} \alpha_{n-1}.$$

The dual basis is

$$\alpha_1 \alpha'_1; \begin{matrix} e_1 & \alpha'_1 \\ \alpha_1 & \alpha'_1 \alpha_1 \end{matrix}; \begin{matrix} e_2 & \alpha'_2 \\ \alpha_2 & \alpha'_2 \alpha_2 \end{matrix}; \dots; \begin{matrix} e_{n-1} & \alpha'_{n-1} \\ \alpha_{n-1} & \alpha'_{n-1} \alpha_{n-1} \end{matrix}; e_n.$$

It is easy to know that $L(A)$ is an ideal of $Z(A)$ generated by $\{\alpha_1 \alpha'_1, \alpha_1 \alpha'_1 + \alpha_2 \alpha'_2, \alpha_2 \alpha'_2 + \alpha_3 \alpha'_3, \dots, \alpha_{n-2} \alpha'_{n-2} + \alpha_{n-1} \alpha'_{n-1}, \alpha'_{n-1} \alpha_{n-1}\}$ and $H(A)$ is generated by $\{2\alpha_1 \alpha'_1 + \alpha_2 \alpha'_2, \alpha_1 \alpha'_1 + 2\alpha_2 \alpha'_2 + \alpha_3 \alpha'_3, \alpha_2 \alpha'_2 + 2\alpha_3 \alpha'_3 + \alpha_4 \alpha'_4, \dots, \alpha_{n-3} \alpha'_{n-3} + 2\alpha_{n-2} \alpha'_{n-2} + \alpha_{n-1} \alpha'_{n-1}, \alpha_{n-2} \alpha'_{n-2} + 2\alpha_{n-1} \alpha'_{n-1}\}$.

Then $\dim_K L(A) = n$ since the rank of the matrix below is n .

$$\begin{bmatrix} 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & 0 & \dots & 0 & 0 \\ & \dots & & \dots & & \dots & \\ 0 & 0 & 0 & 0 & \dots & 1 & 1 \\ 0 & 0 & 0 & 0 & \dots & 0 & 1 \end{bmatrix}_{(n+1) \times n}$$

We know that $\dim_K H(A) < n$ if $\text{Char} K$ is a factor of $n+1$ and $\dim_K H(A) = n$ otherwise, since the determinant of the matrix below is $n+1$.

$$\begin{bmatrix} 2 & 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 2 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 2 & 1 & \dots & 0 & 0 \\ & \dots & & \dots & & \dots & \\ 0 & 0 & 0 & 0 & \dots & 2 & 1 \\ 0 & 0 & 0 & 0 & \dots & 1 & 2 \end{bmatrix}_{n \times n}$$

Then $H(A) \subsetneq L(A)$ if $\text{Char} K$ is a factor of $n+1$ and $H(A) = L(A)$ otherwise.

Example Let K be a field and $A = K[x]/(x^n)$, where $n \in \mathbb{N}$. Clearly, A is a symmetric cellular algebra with a basis $1, \bar{x}, \dots, \overline{x^{n-1}}$. It is easy to know that $L(A)$ has a basis $\overline{x^{n-1}}$ and $Z(A) = A$. This example shows that $\dim_K Z(A) - \dim_K L(A)$ may be very large.

Proposition 3.4. *Notations are as in Theorem A. Then $x_\lambda x_\mu = 0$ for arbitrary $\lambda, \mu \in \Lambda$ with $\lambda \neq \mu$.*

Proof: For arbitrary $\lambda, \mu \in \Lambda$ with $x_\lambda x_\mu \neq 0$, then there exist $S_0 \in M(\lambda)$ and $U_0 \in M(\mu)$ such that

$$C_{S_0, T}^\lambda D_{S_0, T}^\lambda C_{U_0, V}^\mu D_{U_0, V}^\mu \neq 0.$$

This implies $D_{S_0, T}^\lambda C_{U_0, V}^\mu \neq 0$. Then by Corollary 3.1, there exists some $C_{X, Y}^\varepsilon$ such that

$$r_{(U_0, V, \mu), (X, Y, \varepsilon), (S_0, T, \lambda)} \neq 0.$$

By (C3) of Definition 2.5, this implies that $\lambda \leq \mu$. By $x_\lambda x_\mu = x_\mu x_\lambda$, we get $x_\lambda x_\mu \neq 0$ also implies that $\mu \leq \lambda$. Then $\lambda = \mu$ if $x_\lambda x_\mu \neq 0$. $\square$

4 Semisimple case

In this section, we consider the semisimple case. We will construct all the central idempotents which are primitive in $Z(A)$ by elements x_λ defined in Section 3 for a semisimple symmetric cellular algebra.

Firstly, let us recall the definition of Schur elements. For details, see [9].

Let R be a commutative ring with identity and A an R -algebra. Let V be an A -module which is finitely generated and free over R . The algebra homomorphism

$$\rho_V : A \rightarrow \text{End}_R(V), \quad \rho_V(a)v = av, \quad \text{where } v \in V, a \in A,$$

is called the representation afforded by V . The corresponding character is the R -linear map defined by

$$\chi_V : A \rightarrow R, \quad a \mapsto \text{tr}(\rho_V(a)),$$

where tr is the usual trace of a matrix.

Let K be a field and A a finite dimensional symmetric K -algebra with symmetrizing trace τ . Let $B = \{B_i \mid i = 1, \dots, n\}$ be a basis and $D = \{D_i \mid i = 1, \dots, n\}$ the dual basis determined by τ . If V is a split simple A -module, denote the character by χ_V , we have

$$\sum_i \chi_V(b_i) \chi_V(D_i) = c_V \dim_K V,$$

where $c_V \in K$ is the so-called Schur element associated with V . We also denote it by c_{χ_V} . It is non-zero if and only if V is a split simple projective A -module [9].

Lemma 4.1. ([9] 7.2.7) *Let A be a split semisimple K -algebra. Then*

$$\{e_V := c_V^{-1} \sum_i \chi_V(b_i) D_i \mid V \text{ is a simple } A\text{-module}\}$$

is a complete set of central idempotents which are primitive in $Z(A)$.

Let R be an integral domain and A a symmetric cellular algebra with cellular basis $\{C_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$. Given a symmetrizing trace τ , the dual basis is $\{D_{S,T}^\lambda \mid S, T \in M(\lambda), \lambda \in \Lambda\}$. Let K be the field of fractions of R . Define $A_K := A \otimes_R K$. Consider A as a subalgebra of A_K and extend τ of A to A_K . Then we can construct all the central idempotents which are primitive in $Z(A_K)$ by x_λ .

Proposition 4.2. *If A_K is split semisimple, then $\{c_{W(\lambda)}^{-1} x_\lambda \mid \lambda \in \Lambda\}$ is a complete set of central idempotents which are primitive in $Z(A_K)$.*

Proof: The left A_K -module $W(\lambda)$ is split simple since A_K is split semisimple. Then by Lemma 4.1, we have

$$e_{W(\lambda)} = c_{W(\lambda)}^{-1} \sum_{\mu \in \Lambda, U, V \in M(\mu)} \chi_{W(\lambda)}(C_{U,V}^\mu) D_{U,V}^\mu.$$

Note that the character afforded by $W(\lambda)$ is given by the following formula

$$\chi_{W(\lambda)}(a) = \sum_{S \in M(\lambda)} r_a(S, S)$$

for all $a \in A$. Then we get $\chi_{W(\lambda)}(C_{U,V}^\mu) = \sum_{S \in M(\lambda)} r_{(U,V,\mu),(S,T,\lambda),(S,T,\lambda)}$. Then

$$\begin{aligned} e_{W(\lambda)} &= c_{W(\lambda)}^{-1} \sum_{\mu \in \Lambda, U, V \in M(\mu)} \sum_{S \in M(\lambda)} r_{(U,V,\mu),(S,T,\lambda),(S,T,\lambda)} D_{U,V}^\mu, \\ &= c_{W(\lambda)}^{-1} \sum_{S \in M(\lambda)} \sum_{\mu \in \Lambda, U, V \in M(\mu)} r_{(U,V,\mu),(S,T,\lambda),(S,T,\lambda)} D_{U,V}^\mu, \\ &= c_{W(\lambda)}^{-1} \sum_{S \in M(\lambda)} C_{S,T}^\lambda D_{S,T}^\lambda. \end{aligned}$$

□

Remark. Clearly, $\{x_\lambda \mid \lambda \in \Lambda\}$ is a basis of the center of A_K by this proposition. It is different from the one in [12] when we consider Hecke algebras.

Note that the center of A is equal to the intersection of A and the center of A_K . We now give a necessary condition for an element of the center of A_K being in A .

Corollary 4.3. *Let $a_\lambda \in K$ for all $\lambda \in \Lambda$ and $a = \sum_{\lambda \in \Lambda} a_\lambda x_\lambda \in A$. Then $a_\lambda c_{W(\lambda)} n_\lambda \in R$ for arbitrary $\lambda \in \Lambda$, where n_λ is the number of elements in the set $M(\lambda)$.*

Proof: For any $\lambda \in \Lambda$, we know $c_{W(\lambda)}^{-1}x_\lambda$ is a central idempotent of A_K by Proposition 4.2, i.e. $x_\lambda^2 = c_{W(\lambda)}x_\lambda$. This implies that $ax_\lambda = a_\lambda c_{W(\lambda)}x_\lambda$. Clearly, $ax_\lambda \in A$ implies $\tau(ax_\lambda) \in R$. By the definition of the dual basis, $\tau(x_\lambda) = m_\lambda$. This completes the proof. $\square$

Acknowledgments The author acknowledges his supervisor Prof. C.C. Xi and the support from the Research Fund of Doctor Program of Higher Education, Ministry of Education of China. He also acknowledges Dr. Wei Hu for many helpful conversations.

References

- [1] C.W. Curtis and I. Reiner, Representation theory of finite groups and associative algebras, Interscience, New York, 1964.
- [2] R. Dipper and G. James, Blocks and idempotents of Hecke algebras of general linear groups, Proc. London Math. Soc. (3), **54**, (1987), 57-82.
- [3] A. Francis, The minimal bases for the centre of an Iwahori-Hecke algebra, J. Algebra, **221**, (1999), 1-28,
- [4] A. Francis, Centralizers of Iwahori-Hecke algebras, Tran. Amer. Math. Soc., **353**, Number 7, (2001), 2725-2739.
- [5] A. Francis and J.J. Graham, Centers of Hecke algebras: The Dipper-James conjecture, J. Algebra, **306**, (2006), 244-267.
- [6] A. Francis and L. Jones, On bases of centres of Iwahori-Hecke algebras of the symmetric group, J. Algebra, **289**, (2005), 42-69.
- [7] M. Geck, Hecke algebras of finite type are cellular, Invent. math., **169**, (2007), 501-517.
- [8] J.J. Graham and G.I. Lehrer, Cellular algebras, Invent. Math., **123**, (1996), 1-34.
- [9] M. Geck and G. Pfeiffer, Characters of finite Coxeter groups and Iwahori-Hecke algebras, Lond. Math. Soc. Monographs, New series, vol. 21, Oxford University Press, New York (2000).

- [10] M. Geck and R. Rouquier, Centers and simple modules for Iwahori-Hecke algebras, in: *Finite Reductive Groups*, Luminy, 1994, Birkhauser Boston, Boston, MA, 1997, 251-72.
- [11] L. Jones, Centers of generic Hecke algebras, *Trans. Amer. Math. Soc.*, **317**, (1990), 361-392.
- [12] Yeon-Kwan Jeong, In-Sok Lee, Hyekyung Oh and Kyung-Hwan Park, Cellular algebras and centers of Hecke algebras, *Bull. Korean Math. Soc.*, **39**, (2002), No.1, 71-79.
- [13] Yanbo Li, Radicals of symmetric cellular algebras, arXiv: 0911.3524v1 [math.RT], preprint (2009).
- [14] H.B. Rui and C.C. Xi, The representation theory of cyclotomic Temperley-Lieb algebras, *Comment. Math. Helv.*, **79**, no.2, (2004), 427-450.
- [15] C.C. Xi, Partition algebras are cellular *Compositio math.*, **119**, (1999), 99-109.
- [16] C.C. Xi, On the quasi-heredity of Birman-Wenzl algebras, *Adv. Math.*, **154**, (2000), 280-298.